arXiv:1202.2798v2 [quant-ph] 2 Mar 2012

Most robust and fragile two-qubit entangled states under depolarizing channels

Chao-Qian Pang,[1] Fu-Lin Zhang,[1,*] Yue Jiang,[1] Mai-Lin Liang,[1] and Jing-Ling Chen[2,3]
[1] *Physics Department, School of Science, Tianjin University, Tianjin 300072, China*
[2] *Theoretical Physics Division, Chern Institute of Mathematics, Nankai University, Tianjin, 300071, China*
[3] *Centre for Quantum Technologies, National University of Singapore, 3 Science Drive 2, Singapore 117543*
(Dated: March 5, 2012)

In the two-qubit system under the local depolarizing channels, the most robust and the most fragile states for a given concurrence or negativity are derived. For the one-sided channel, with the aid of the evolution equation for entanglement given by Konrad *et al.* [Nat. Phys. 4, 99 (2008)], the pure states are proved to be the most robust. Based on a generalization of the evolution equation, we classify the ansatz states in our investigation by the amount of robustness, and consequently derive the most fragile states. For the two-sided channel, the pure states are proved to be the most robust for a fixed concurrence, but is the most fragile with a given negativity when the channel is uniform. Under the uniform channel, for a given negativity, the most robust states are the ones with the maximal concurrence, which are also the most fragile states when the concurrence is given in the region of $[1/2, 1]$. When the entanglement approaches zero, the most fragile states for a given negativity become the pure states, but the ones for concurrence turn into a mixture of two separable pure states.

PACS numbers: 03.67.Mn,03.65.Ud,03.65.Yz

I. INTRODUCTION

Quantum coherent superposition makes quantum information conceptually more powerful than classical information, which is the essential distinction between a quantum system and a classical one. Entanglement is a manifestation of the distinction in composite systems [1] and is one of the key resources in the field of quantum information science [2]. However, the unavoidable couplings between a real quantum system and its environment can lead to the decoherence phenomenon. Simultaneously, the entanglement between the subsystems is usually destroyed.

Because of the importance both in the fundamental theory and the application in quantum information, the dynamics of entanglement in a quantum system under decoherence has attracted wide attention in recent years. Many significant and interesting results have been reported. For instance, the concept of sudden death of entanglement (ESD) has been presented by Yu and Eberly [3, 4], which means that the entanglement can decay to zero abruptly in a finite time while complete decoherence requires an infinite amount of time. This interesting phenomenon has been recently observed in two sophistically designed experiments with photonic qubits [5] and atomic band [6]. In [7], by utilizing the Jamiołkowski isomorphism, Konrad *et al.* presented a factorization law for a two qubit system, which described the evolution of entanglement in a simple and general way. This result has been extended in many directions [8–11].

This paper is concerned about the topic of the robustness of entanglement, when a quantum system interaction with a noise environment. Vidal and Tarrach [12] introduced the robustness of entanglement as a measure of entanglement, corresponding to the minimal amount of mixing with separable states which washed out all entanglement. Subsequently, Simon and Kempe [13] considered the critical amount of depolarization, where the entanglement vanishes, as a quantitative signature of the robustness of the entanglement, when they studied the robustness of multi-party entanglement under local decoherence, modeled by partially depolarizing channels acting independently on each subsystem. This definition was adopted in the recent work [14], in which an interesting residual effect was pointed out, on the robustness of a three-qubit system in an arbitrary superposition of Greenberger-Horne-Zeilinger state and W state.

We notice that, even in the two-qubit system, there are still some questions about the robustness to explore. For instance, the results of [14] show the robustness increases synchronously with the degree of entanglement for a two-qubit pure state. Obviously, it should be very hard to extend this conclusion to the mixed states, because the measurements of entanglement for the mixed states are not unique [15–17], which would influence the robustness jointly . Then, a question arises: For a given value of entanglement under a measurement, which states are the most robust and which are the most fragile? Since, generally speaking, the value of entanglement in a two-qubit system correlated positively with the resource of the preparation [15, 16] and its capacity in quantum information [18, 19], the entanglement is expected to be stable under the same conditions. Many similar investigations have been reported for about ten years. For example, the famous *maximally entangled mixed states* [20, 21] can be considered to be the most residual entanglement under a global noise channel. In [22], Yu and Eberly studied the robustness and fragility of entangle-

---

[*] Email: flzhang@tju.edu.cn

ment in some exactly solvable dephasing models. In the recent work of Novotný *et al.* [23], they present a general and analytically solvable decoherence model without any weak-coupling or Markovian assumption, and distinguish the robust and fragile states in the model.

In the present work, we investigate the two-qubit system under local depolarizing channels which is invariant under local unitary operations, and focus on the most robust entangled states (MRES) and the most fragile entangled states (MFES) for a given amount of initial entanglement. The model in [13] and [14] is generalized to the nonuniform case in our study. Namely, the strength of the coupling of the two qubits with its environment are different. This generalization is more close to some practical cases , in which the entangled two qubits undergo different environment. Two measurements of the degree of entanglement, the concurrence [15, 16] and the negativity [17], are adopted for comparison. In Sec. II, we review the definitions and introduce some notations in this paper. Considering its relation with the *evolution equation for quantum entanglement* in [7], we selectively give the result of the one-sided channel in Sec. III, which is an extreme case of the nonuniform channels. In Sec. IV, the uniform and nonuniform channels are studied together. Conclusion and discussion are made in the last section.

II. NOTATIONS AND DEFINITIONS

A. Channels and robustness

Under local decoherence channels, without the interaction between the two qubits, the dynamics of each qubit is governed by a master equation which depends on its own environment [24]. The evolution of each reduced density matrices is described by a completely positive trace-preserving map: $\rho_i(t) = \varepsilon_i(t)\rho_i(0)$, $i = 1, 2$, and for the whole state $\rho(t) = \varepsilon_1(t) \otimes \varepsilon_2(t)\rho(0)$. In the Born-Markov approximation these maps (or channels) can be described using its Kraus representation

$$\varepsilon_i(t)\rho_i(0) = \sum_{j=0}^{M} E_{ji}\rho_i(0)E_{ji}^{\dagger} \tag{1}$$

where E_{ji} satisfying $\sum_{j=0}^{M-1} E_{ji}E_{ji}^{\dagger} = I$, are the Kraus operators, M being the number of operators needed to completely characterize the channel. For the depolarizing channels, the Kraus operators are

$$E_{0i} = \frac{1}{2}\sqrt{3s_i + 1}I_i, \quad E_{ji} = \frac{1}{2}\sqrt{1 - s_i}\sigma_{ji}, \tag{2}$$

where σ_{ji}, $(j = 1, 2, 3)$ is the jth Pauli matrices for the ith qubit. We consider the noise parameter $s_i = \exp(-\kappa_i t)$, where κ_i is the decay constant determined by the strength of the interaction of the ith qubit with its environment and t is the interaction time [25]. In the Bloch sphere representation for the qubit density operator, the transformation in Eqs. (1) and (2) is given by $\rho_i(t) = \frac{1}{2}(I_i + s_i\vec{r}_i \cdot \vec{\sigma}_i)$ with $\vec{r}_i$ being the the initial Bloch vector.

For simplicity but without loss of generality, we choose the decay constants $\kappa_1 = 1 + \Delta$ and $\kappa_2 = 1 - \Delta$, with the uniform parameters $\Delta \in [0, 1]$. Then, the noise parameters can be written as $s_1 = s^{1+\Delta}$ and $s_2 = s^{1-\Delta}$, where $s = e^{-t}$. Here, we introduce the measurement of the robustness for a state ρ under a nonuniform channel as

$$\mathcal{R}(\rho) = 1 - s_{crit}(\rho), \tag{3}$$

where $s_{crit}(\rho)$ denotes the critical value of the noise parameter s, at which occurs the ESD of the two-qubit system with the initial state ρ. When $\Delta = 0$, it returns the definition in [13] and [14].

B. Measurements of entanglement

The concurrence of a pure two-qubit state $|\psi\rangle = c_1|00\rangle + c_2|01\rangle + c_3|10\rangle + c_4|11\rangle$ is given by $\mathcal{C}(|\psi\rangle) = 2|c_1c_4 - c_2c_3|$. For a mixed state, the concurrence is defined as the average concurrence of the pure states of the decomposition, minimized over all decompositions of $\rho = \sum_j p_j|\psi_j\rangle\langle\psi_j|$,

$$\mathcal{C}(\rho) = \min \sum_j p_j\mathcal{C}(|\psi_j\rangle). \tag{4}$$

It can be expressed explicitly as [15, 16] $\mathcal{C}(\rho) = \max\{0, \sqrt{\lambda_1} - \sqrt{\lambda_2} - \sqrt{\lambda_3} - \sqrt{\lambda_4}\}$, in which $\lambda_1, ..., \lambda_4$ are the eigenvalues of the operator $R = \rho(\sigma_y \otimes \sigma_y)\rho^*(\sigma_y \otimes \sigma_y)$ in decreasing order and σ_y is the second Pauli matrix.

For a bipartite system described by the density matrix ρ, the negativity is defined as [17, 26]

$$\mathcal{N}(\rho) = 2\sum_j |\lambda_j|, \tag{5}$$

where λ_j are the negative eigenvalues of ρ^{T_2} and T_2 denotes the partial transpose with respect to the second subsystem. For the two-qubit system, the partial transpose density matrix ρ^{T_2} has at most one negative eigenvalue [21].

C. Ansatz states

To derive the MRES and MFES for a given value of concurrence or negativity, we adopt the approach in [20, 21]. Under a given value of the nonuniform parameter Δ, we randomly generate two-qubit states and obtain their robustness and degree of entanglement by numerical computation. Plotting them in the corresponding robustness-entanglement planes (such as the ones in Fig.

1), we fortunately find, for any value of Δ, their regions are always the same as the states

$$\rho_{ansatz}(r,\theta) = r|\psi(\theta)\rangle\langle\psi(\theta)| + (1-r)|01\rangle\langle 01|, \qquad (6)$$

where $r \in [0,1]$ and $|\psi(\theta)\rangle = \cos\theta|00\rangle + \sin\theta|11\rangle$ with $\theta \in [0, \pi/2]$. For each value of $\Delta \in \{0, 0.1, 0.2, ...1\}$, and for both the entanglement measures, this is supported by 300000 random states. Therefore, the MRES or MFES can be represented as the state (6) with a constraint, which we will derive in the following sections. We call the state (6) the *ansatz state* in this paper.

One thing to be mentioned is that, The two extreme cases of the states are corresponding to, the pure states with the same values of negativity and concurrence when $r = 1$, and the ones with the minimal negativity for a given concurrence when $\theta = \pi/4$, respectively [27].

III. ONE-SIDED NOISY CHANNEL

A. Robustness vs concurrence

When $\Delta = 1$, $s_1 = s^2$ and $s_2 = 1$, the channel reduces the one-sided depolarizing channel , in which only the first qubit influenced by the environment. The central result of [7] is the factorization law for an arbitrary pure state $|\psi\rangle$ under a one-sided noisy channel

$$\mathcal{C}[(\$ \otimes I)|\psi\rangle\langle\psi|] = \mathcal{C}[(\$ \otimes I)|\psi^+\rangle\langle\psi^+|]\mathcal{C}(|\psi\rangle), \qquad (7)$$

where $|\psi^+\rangle = |\psi(\pi/4)\rangle$ and \$ denotes an arbitrary channel not restricted to a completely positive trace-preserving map. An evident corollary is that, the ESD of the systems initial from any entangled pure states occur at the same time, and consequently their robustness are the same.

An application of (7) in [7] is the inequality for mixed states ρ_0,

$$\mathcal{C}[(\$ \otimes I)\rho_0] \leq \mathcal{C}[(\$ \otimes I)|\psi^+\rangle\langle\psi^+|]\mathcal{C}(\rho_0). \qquad (8)$$

From (7) and (8), one can find the fact that, under the same one-sided channel, with the same initial concurrence, the ESD of a mixed state comes not later than the one of a pure state. Hence, the MRES with a fixed concurrence is the pure sates. Under the depolarizing channel considered in the present work, we can derive the value of robustness

$$\mathcal{R}_{MRES} = \mathcal{R}(|\psi\rangle) = 1 - \frac{1}{\sqrt{3}}, \qquad (9)$$

which is a constant, independent of the concurrence.

To obtain the MFES, we generalize the factorization law (7) in the arbitrary two-qubit states case. Namely, if two states satisfy

$$\rho_1 = \gamma(I \otimes M)\rho_0(I \otimes M^\dagger) \qquad (10)$$

with $1/\gamma = \mathrm{Tr}[(I \otimes M)\rho_0(I \otimes M^\dagger)]$ and $M = I + \vec{a}\cdot\vec{\sigma}$, ($a \in [0,1]$), the evolutions of their concurrence are related as

$$\mathcal{C}[(\$ \otimes I)\rho_1]\mathcal{C}(\rho_0) = \mathcal{C}[(\$ \otimes I)\rho_0]\mathcal{C}(\rho_1). \qquad (11)$$



FIG. 1: (color online) Plots of the states $\rho(c,p)$ in the (a) robustness-concurrence and (b) robustness-negativity planes under one-sided noise channel. The dotted lines correspond to the parameter $c = 1, 0.8, 0.6, 0.4, 0.2$ from top to bottom, in both (a) and (b). The dashed lines refer to the parameter (1) $p = \tan(0.1 \times \pi/2)$, (2) $p = \tan(0.2 \times \pi/2)$, (3) $p = \tan(0.3 \times \pi/2)$ and (4) $p = \tan(0.4 \times \pi/2)$. The solid line is the MFES in the two cases.

We don't give the proof of the above relation, which is exactly the same as the process for (7) in [7]. As the case of the pure states, we can conclude that the robustness of ρ_0 and ρ_1 in (10) have the same value.

Then, we find the ansatz states (6) can be classified by the amount of robustness, with the aid of the generalized factorization law (11). Namely, choosing $\rho_0 = \rho_{ansatz}(c, \pi/4)$ and $\vec{a} = (0, 0, (1-p)/(1+p))$, the ansatz states can be written as

$$\rho_{ansatz}(r,\theta) = \rho(c,p) = \gamma(I \otimes M)\rho_0(I \otimes M^\dagger), \qquad (12)$$

where the relations between the two pairs parameters are

$$c = \frac{r - r\cos 2\theta}{1 - r\cos 2\theta}, \qquad p = \tan\theta. \qquad (13)$$

The robustness of the ansatz state is only dependent of the parameter c in ρ_0, which is given by

$$\mathcal{R}[\rho(c,p)] = 1 - \sqrt{\frac{2-c}{2+c}}, \qquad (14)$$

corresponding the result for the pure states in (9) when

$c=1$. The concurrence can be obtained as

$$\mathcal{C}[\rho(c,p)] = \frac{2cp}{c+(2-c)p^2} = r\sin 2\theta. \qquad (15)$$

In Fig. 1 (a), we plot the ansatz states with some discrete values of c or p in the robustness-concurrence plane. One can find the MFES, with the minimal robustness for a given concurrence, achieves the maximal concurrence when the value of robustness is fixed. It is easy to determine the maximal value of concurrence in (15) with a fixed c. Then the constraint on the ansatz sates for the MFES can be obtained as

$$c(1+p^2) - 2p^2 = 0, \qquad (16)$$

with $0 \le p \le 1$, or equivalently

$$2r\cos^2\theta = 1, \qquad (17)$$

with $0 \le \theta \le \pi/4$ and $1/2 \le r \le 1$. It is interesting that the condition leads to the lengthes of the Bloch vectors for the two qubit are $r_1 = 2(1-r)$ and $r_2 = 0$, which reach the maximal difference with a fixed r.

B. Robustness vs negativity

It is easily to prove the the fact that, the MRES under an arbitrary one-sided noise channel for a given negativity are also the pure states, with the aid of the results for the concurrence case. Suppose ρ denotes an arbitrary entangled two-qubit state, $|\psi_1\rangle$ and $|\psi_2\rangle$ are two pure states, and $\mathcal{N}(\rho) = \mathcal{N}(|\psi_1\rangle)$ and $\mathcal{C}(\rho) = \mathcal{C}(|\psi_2\rangle)$. Two relations can be obtained in the above subsection as $\mathcal{R}(\rho) \le \mathcal{R}(|\psi_2\rangle)$ and $\mathcal{R}(|\psi_1\rangle) = \mathcal{R}(|\psi_2\rangle)$, from which we deduce the conclusion.

In Fig. 1 (b), one can notice the region of the ansatz states, of course the one of the arbitrary states, in the robustness-negativity plane is similar to the one in Fig. 1 (a) where the entanglement measured by concurrence. Thus, to determine the MFES is to find the states $\rho(c,p)$ with the maximal negativity for a fixed c. The negativity of the ansatz states is

$$\mathcal{N}[\rho_{ansatz}(r,\theta)] = \sqrt{r^2\sin^2 2\theta + (1-r)^2} - (1-r). \qquad (18)$$

Inserting the relations (13), one can find the maximal negativity occurs when

$$p = \sqrt{\frac{2\sqrt{2-c}(-1+c)c^{3/2} + 2c^2 - c^3}{-8+24c-20c^2+5c^3}}. \qquad (19)$$

The corresponding relation of the parameters r and θ is very close to the numerical results for $\Delta = 0.99$ in Fig. 5 (a) in the following section.

In the MFES, when the parameter $\theta \to 0$, the entanglement of the entangled composition decreases, its proportion $r \to 1$. However, in the MFES for concurrence given in (16) and (17), when $\theta \to 0$, the proportion r drops to $1/2$, which has the maximal entropy in the family of the ansatz states.



FIG. 2: (color online) The relation between the robustness and the concurrence of a two-qubit system in a pure entangled state. From top to bottom the curves refer to $\Delta = 1, 0.75, 0.5, 0.25, 0$.

IV. TWO-SIDED NOISY CHANNEL

We fail in generalizing the relations in (10) and (11) to the two-sided channel case. Therefore, we calculate the ESD critical condition of the ansatz states directly

$$\begin{aligned}\mathcal{P} = 4r^2\sin^2 2\theta s_1^2 s_2^2 - [1+(1-2r)s_1 s_2]^2 \\ +[r\cos 2\theta(s_1-s_2) + (1-r)(s_1+s_2)]^2 = 0,\end{aligned} \qquad (20)$$

which is one of the main basis to determine the MRES and MFES in this section. When $r=1$, one can obtain the relation of the entanglement of pure states and their critical noise parameters

$$\mathcal{C}^2(|\psi\rangle) = \mathcal{N}^2(|\psi\rangle) = \frac{(1-s_1^2)(1-s_2^2)}{4s_1^2 s_2^2 - (s_1-s_2)^2}. \qquad (21)$$

For the Bell state $|\psi^+\rangle$, the above equation gives the robustness $\mathcal{R}(|\psi^+\rangle) = 1 - 1/\sqrt{3}$, which is independent of the nonuniform parameter Δ. In this sense, our definition of robustness is reasonable. When $\Delta = 0$, $\mathcal{R}(|\psi\rangle) = 1 - 1/\sqrt{2\mathcal{C}(|\psi\rangle)+1}$, which is the result of the uniform channels given in [14]. In Fig. 2, we plot the relation of the robustness and the entanglement of the pure state. Under a fixed Δ, the robustness is a monotone increasing function of the entanglement. A nonmaximal entangled pure state becomes more fragile as Δ decreases.

A. Robustness vs concurrence

Because of the invariance of the depolarizing channels under local unitary (LU) transformations, the MRES for concurrence can be proved to be the pure states under the two-sided channels. According to the procedure given by Wootters [16], one can always obtain a decomposition $\{|\phi_i\rangle\}$ minimizing the average concurrence in Eq. (4),



FIG. 3: (color online) The relation between the robustness and the concurrence of the MRES and MFES under different nonuniform parameters, $\Delta = 0.75, 0.5, 0.25, 0$. The dots on the lines denote the quasi-MFES, whose parameters $\beta = 0.1, 0.2, 0.3, 0.4$, with the values of concurrence increasing .

$\rho_0 = \sum_i t_i |\phi_i\rangle\langle\phi_i|$, in which $\sum_i t_i = 1$ and all the elements have the same value of concurrence as the mixed state ρ_0. The elements are equivalent under LU transformation to the same state $|\psi\rangle$

$$|\phi_i\rangle \ = U_i^A \otimes U_i^B |\psi\rangle, \tag{22}$$

with the concurrence $\mathcal{C}(|\psi\rangle) = \mathcal{C}(\rho_0)$. After passage through a basis independent local channel $\$_1 \otimes \$_2$, the concurrence $\mathcal{C}[(\$_1 \otimes \$_2)|\phi_i\rangle\langle\phi_i|] = \mathcal{C}[(\$_1 \otimes \$_2)|\psi\rangle\langle\psi|]$. It then immediately follows, by convexity, that

$$\mathcal{C}[(\$_1 \otimes \$_2)\rho_0] \leq \mathcal{C}[(\$_1 \otimes \$_2)|\psi\rangle\langle\psi|], \tag{23}$$

which leaves us with the claim at the beginning of this paragraph.

Similar with the case of the one-sided depolarizing channels, the MFES in the present case are also the states achieve the maximal concurrence for a given value of robustness, which can be noticed in Fig. 3. From (15) and (20), we find, for a given pair of the critical noise paraments s_1 and s_2, or equivalently the robustness $\mathcal{R}$ and the nonuniform parameter Δ, the MFES can be obtained applying the restraint

$$\alpha = \frac{1}{4}\left[1 + \sqrt{8\Omega(2\beta - 1)\beta + 1}\right] \tag{24}$$

on the ansatz states (6), where $\alpha = r\cos^2\theta \in [0, 1/2]$, $\beta = r\sin^2\theta \in [0, 1/2]$ and $\Omega = \frac{s_1^2(1-s_2^2)}{s_2^2(1-s_1^2)}$. When $\Delta = 1$, $\Omega = 0$, the restraint, $\alpha = 1/2$, returns to the result of the one-sided channels in (16) and (17).

For the uniform noise channels, where $\Delta = 0$ and $\Omega = 1$, the MFES can be determined as

$$\rho_{MFES} = \begin{cases} \frac{1}{2}[|\psi(\theta)\rangle\langle\psi(\theta)| + |01\rangle\langle 01|], & \mathcal{C} < 1/2\,, \\ r|\psi^+\rangle\langle\psi^+| + (1-r)|01\rangle\langle 01|, & \mathcal{C} \geq 1/2\,, \end{cases} \tag{25}$$

by using the relation in (24). In the first region where $\mathcal{C} < 1/2$, the MFES reach the maximal entropy in the family of the ansatz states. In the other, when $\mathcal{C} \geq 1/2$, the MFES are the states with the minimal negativity for a fixed concurrence [27].

When $0 < \Delta < 1$, the MFES for the arbitrary depolarizing channels is given by the simultaneous equations of (20) and (24), which is shown in Fig. 3 for some values of Δ. It can be noticed that, the difference between the robustness of MRES and MFES decreases with the decrease in the nonuniform parameter Δ. When $\beta \to 0$, the concurrence $\mathcal{C} \to 0$, the proportion of the entangled state in MFES approaches 1/2.

Considering the absence of a brief expression of the MFES, we present a family of quasi-MFES. Namely, we replace the critical depolarizing parameter $s^2 \in [1/3, 1]$ in (24) by its mean value 2/3, and obtain $\Omega \to \widetilde{\Omega} = \frac{(2/3)^{\Delta-1}-1}{(2/3)^{-\Delta-1}-1}$. It is obviously, $\widetilde{\Omega}|_{\Delta=0} = 1$ and $\widetilde{\Omega}|_{\Delta=1} = 0$.The positions of the quasi-MFES in the robustness-concurrence plane are shown in Fig. (3), which are very close to the MFES. Our numerical result shows, for a given β, the fidelity of the quasi-MFES and the MFES $F(\rho_{quasi-MFES}, \rho_{MFES}) \geq 1 - 10^{-4}$ with $F(\rho_1, \rho_2) = \left[\mathrm{Tr}\left(\sqrt{\sqrt{\rho_1}\rho_2\sqrt{\rho_1}}\right)\right]^2$, which is supported by one million pairs of randomly generated quasi-MFES and MFES.

B. Robustness vs negativity

To determine the MRES and MFES for the negativity under the two-sided depolarizing channels, we derive the condition,

$$\frac{\partial\mathcal{N}}{\partial r}\frac{\partial\mathcal{P}}{\partial\theta} - \frac{\partial\mathcal{N}}{\partial\theta}\frac{\partial\mathcal{P}}{\partial r} = 0, \tag{26}$$

where the negativity of an ansatz state reaches an extremum for a fixed robustness. The form of the the MRES and MFES are given by the physically accepted solutions to the simultaneous equations of (20) and (26).

When $\Delta = 0$, one can check the two solutions $\theta = \pi/4$ and $r = 1$, corresponding to the MRES and MFES respectively. In other words, under the uniform channel, the MRES for the entanglement measurement negativity are $\rho_{MRES} = \rho_{ansatz}(r, \pi/4)$, and the MFES $\rho_{MFES} = \rho_{ansatz}(1, \theta)$ are the pure states. The result is contrary to the concurrence case, where the pure states are the most robust, and $\rho_{ansatz}(r, \pi/4)$ are the most fragile (when $\mathcal{C} \geq 1/2$). The difference indicate the robustness is influenced by both of the two measures of entanglement. The pure states, which are the MRES for concurrence, have the maximal negativity when the concurrence is fixed. They are also the states with a minimal concurrence for a given negativity, and thus are the MFES for negativity. The states $\rho_{ansatz}(r, \pi/4)$ are the states having the minimal negativity for a fixed concurrence, and also the states with maximal concurrence for a given negativity. Therefore, they appear to be more fragile than the other states with the same concurrence, but more robust than the other states with the same neg-



FIG. 4: (color online) Plots of the states $\rho(r,\theta)$ in the robustness-negativity planes with the MRES and MFES when the nonuniform parameters $\Delta = 0.5$. The parameters r/θ takes a set of discrete values in (a)/(b), which are labeled nearby the tangent points with the MRES/MFES.



FIG. 5: (color online) Relations of the parameters r and θ for the MFES (a) and MRES (b) with different inhomogeneous parameters, $\Delta = 0.99, 0.75, 0.5, 0.25, 0$. The robustness takes a set of regularly spaced values.

ativity.

It is impossible to derive an explicit expression of the solution to (20) and (26) for the arbitrary $\Delta \in (0,1)$. Thus we plot the ansatz states in the robustness-negativity plane for various values of the nonuniform parameter Δ, one of which is shown in Fig. 4. One can notice that, the MRES can be expressed as the ansatz states with a constrain as $\theta = \theta(r)$, with r varying from 0 to 1. And the constrain for MRES is $r = r(\theta)$, where the domain is $\theta \in [0, \pi/4]$. Then we search the constrains by numerical computation, and plot them in Fig. 5. Obviously, the proportion of the entangled composition r in the MFES is always more than 3/4. The proportion approaches 1, when the entanglement $\mathcal{N} \to 0$. It increases with decrease in the nonuniform parameter Δ. Under a fixed Δ, the parameter θ alters little when the robustness varies in a vary large region. It decreases to $\pi/4$ quickly, when the r approaches 1, and the robustness approaches the maximum $1-1/\sqrt{3}$. When $\Delta \to 1$, the domain of the robustness becomes a point $\mathcal{R} = 1 - 1/\sqrt{3}$, where $r = 1$ and θ varies from $\pi/2$ to $\pi/4$.

V. CONCLUSION AND DISCUSSION

In conclusion, we investigate the robustness of the entangled states in the two-qubit system, under the local depolarizing channels. The MRES and MFES are derived for a given amount of entanglement, measured by concurrence and negativity. Withe a numerical simulation, we find a family of ansatz states (6), which contain the MRES and MFES. The ansatz state is a mixture of two mutually orthogonal pure states, one of which is entangled and the other separable.

Under the one-sided channels, the pure states are proved to be the MRES, for both the two entanglement measurements, by utilizing the factorization law for the evolution of concurrence given by [7]. A generalized factorization law is presented in (10) and (11). Based on this result, we classify the ansatz states by the values of robustness and derive the MRES for the two measures of entanglement. In the MRES for concurrence, the length of the Bloch vector for the first qubit under the noise reached its maximum, and the one for the free qubit is zero.

Under the two-sided channels, for a given concurrence, the pure states are prove to be the most robust. For the uniform channel, the MFES are the states with the minimal negativity when $\mathcal{C} \geq 1/2$, but the ansatz states with the maximal entropy when $\mathcal{C} < 1/2$. The proportion of the entangled pure state the MFES approaches 1/2, when the concurrence $\mathcal{C} \to 0$. In contrast, the MFES for the negativity under the same channels become a separable pure state when the entanglement approaches zero. In the uniform channel, the pure states are the most fragile for a fixed negativity. And the ansatz states with $\theta = \pi/4$ are more robust than the other states with the same negativity. The reason to the difference between the results for concurrence and negativity is the fact that, the robustness is affected by both of the two entanglement measurements. In addition to the concurrence and the negativity, the degree of mixture is a possible factor affecting the robustness, as shown by the MFES in the uniform channel a given concurrence in $[0, 1/2]$. The local properties, such as the Bloch vectors of each qubit, present a significant affection on the robustness when the channel becomes nonuniform.

Based the classification of the ansatz states under one-sided channels, it is easy to study the relation of the robustness and entanglement. Then, can the arbitrary two-qubit states be classified by robustness? The solution to this question would leave us an explicit expression of robustness under one-sided channel. The evolution equations of entanglement given in [7] have been generalized in many directions [8–11]. It is interesting to study robustness of entanglement in for multi-qubit or multi-qudit systems, as an application of the generalized evolution equations. On the other hand, the capacity of an entangled qubit pair in quantum information [18, 19] does not only depend on the amount of entanglement. It is necessary to investigate the robustness of the capacities in a specific quantum information process. Our scenario to derive the MRES or MFES when an analytic proof is absence can be applied directly to these topics.

Acknowledgments

F.L.Z. is supported by NSF of China (Grant No. 11105097). J.L.C. is supported by National Basic Research Program (973 Program) of China under Grant No. 2012CB921900, NSF of China (Grant Nos. 10975075 and 11175089) and also partly supported by National Research Foundation and Ministry of Education, Singapore (Grant No. WBS: R-710-000-008-271).

---

[1] A. Einstein, B. Podosky, and N. Rosen, Physical Review **47**, 777 (1935).
[2] M. A. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, 2000).
[3] T. Yu and J. Eberly, Physical review letters **93**, 140404 (2004).
[4] T. Yu and J. H. Eberly, Physical review letters **97**, 140403 (2006).
[5] M. Almeida, F. De Melo, M. Hor-Meyll, A. Salles, S. Walborn, P. Ribeiro, and L. Davidovich, Science **316**, 579 (2007).
[6] J. Laurat, K. Choi, H. Deng, C. Chou, and H. Kimble, Physical review letters **99**, 180504 (2007).
[7] T. Konrad, F. De Melo, M. Tiersch, C. Kasztelan, A. Aragão, and A. Buchleitner, Nature physics **4**, 99 (2007).
[8] C. Yu, X. X. Yi, and H. S. Song, Physical Review A **78**, 062330 (2008).
[9] Z. G. Li, S. M. Fei, Z. D. Wang, and W. M. Liu, Physical Review A **79**, 024303 (2009).
[10] Z. Liu and H. Fan, Physical Review A **79**, 032306 (2009).
[11] M. Tiersch, F. De Melo, and A. Buchleitner, Physical review letters **101**, 170502 (2008).
[12] G. Vidal and R. Tarrach, Phys. Rev. A **59**, 141 (1999).
[13] C. Simon and J. Kempe, Phys. Rev. A **65**, 052327 (2002).
[14] B.-K. Zhao and F.-G. Deng, Phys. Rev. A **82**, 014301 (2010).
[15] S. Hill and W. K. Wootters, Physical Review Letters **78**, 5022 (1997).
[16] W. K. Wootters, Physical Review Letters **80**, 2245 (1998).
[17] K. Życzkowski, P. Horodecki, A. Sanpera, and M. Lewenstein, Physical Review A **58**, 883 (1998).
[18] J. Lee and M. S. Kim, Physical Review Letters **84**, 4236 (2000).
[19] G. Bowen and S. Bose, Physical Review Letters **87**, 267901 (2001).
[20] W. J. Munro, D. F. V. James, A. G. White, and P. G. Kwiat, Physical Review A **64**, 030302(R) (2001).
[21] T.-C. Wei, K. Nemoto, P. M. Goldbart, P. G. Kwiat, W. J. Munro, and F. Verstraete, Physical Review A **67**, 022110 (2003).
[22] T. Yu and J. H. Eberly, Physical Review B **66**, 193306 (2002).
[23] J. Novotnŷ, G. Alber, and I. Jex, Physical Review Letters **107**, 90501 (2011).
[24] A. Borras, A. P. Majtey, A. R. Plastino, M. Casas, and A. Plastino, Physical Review A **79**, 022108 (2009).
[25] W. Dür and H. Briegel, Physical review letters **92**, 180403 (2004).
[26] G. Vidal and R. F. Werner, Physical Review A **65**, 032314 (2002).
[27] F. Verstraete, K. Audenaert, J. Dehaene, and B. Moor, Journal of Physics A: Mathematical and General **34**, 10327 (2001).